

POMERON LOOPS & DUALITY IN HIGH ENERGY QCD

Edmond IANCU

Service de Physique Théorique, CEA/DSM/SPhT, CE Saclay, F-91191 Gif-sur-Yvette, France

I discuss the physical picture underlying the evolution equations with Pomeron loops recently derived in multicolor QCD at high energy and qualitatively explain the notion of ‘self-duality’.

The rapid rise of the gluon density in a hadronic wavefunction with increasing energy together with the phenomenon of gluon saturation open the way towards a perturbative treatment of high-energy scattering in QCD in the vicinity of the unitarity limit¹. To that aim, one needs evolution equations which include all the microscopic processes contributing in perturbative QCD to the growth of the gluon distribution and its eventual saturation. These processes have been discussed in a different article in these Proceedings² and are also summarized in Fig. 1. They include the *BFKL evolution* (Fig. 1.b), which involves a $2 \rightarrow 2$ gluon vertex and leads to the rapid growth of the gluon density with increasing energy, together with *gluon splitting* vertices $2 \rightarrow n$ with $n > 2$ (Fig. 1.c) and *gluon recombination* vertices $n \rightarrow 2$ (Fig. 1.d). One must emphasize here that *all* such processes — in fact, all the gluon number changing vertices $m \rightarrow n$ with arbitrary $m, n \geq 2$ — are important for computing scattering amplitudes at very high energy³, and this even when the projectile is a relatively simple object, like a *color dipole* (the interesting case for deep inelastic scattering at small x). Indeed, the non-linear effects responsible for gluon saturation and unitarization involve higher n -point correlation functions (cf. Fig. 1.d), which in turn are generated through splitting processes in the low density regime (cf. Fig. 1.c). In Fig. 1, we have also indicated some simple scattering processes involving one or two external dipoles that we shall use below to probe the gluon evolution in the target.

Gluon-number changing vertices started to be computed in perturbative QCD in the early nineties⁴, but their systematic inclusion into evolution equations turned out to be a complicated problem, whose general solution is not yet known. The *merging* vertices are presently included in the functional JIMWLK equation¹, whereas the *splitting* vertices are rather described by the *dual* version of the JIMWLK Hamiltonian⁵ (see below for the notion of ‘duality’). But the complete evolution Hamiltonian, which should include both splitting and mergings, and therefore be *self-dual*^{5,6}, is not yet fully known (see however Refs.⁷).

The desiderata of including both splitting and merging has been nevertheless achieved^{3,8} within the multicolor limit $N_c \rightarrow \infty$, which allows for important technical simplifications and also for the use of a suggestive dipole language^{9,10} (see also Refs.^{6,11,12}).

Namely, at large N_c and in the dilute regime, the gluons in the target wavefunction can be effectively replaced by color dipoles, which interact with the external dipoles from the projectile via two gluon exchanges (compare in this respect the processes in Figs. 1.a and 2.a). Then the BFKL evolution together with the $2 \rightarrow 4$ gluon splitting^a (as previously depicted in Figs. 1.b

^aAt large N_c , and for a dipole projectile, one can ignore all the splitting vertices except for the vertex $2 \rightarrow 4$.




Figure 1: *Elementary processes which occur in one-step evolution in QCD at high energy: (a) tree-level process; (b) standard BFKL evolution; (c) $2 \rightarrow 4$ gluon splitting; (d) $4 \rightarrow 2$ gluon merging.*




Figure 2: *Target evolution in the dipole picture.*

and c) can be represented as one target dipole splitting into two, followed by the interaction between the child dipoles and the projectile (cf. Figs. 2.b and c). Note that, whereas the scattering with a single external dipole serves as a measure of the *average dipole number density* in the target, and thus of the standard BFKL evolution (cf. Fig. 2.b), on the other hand, the scattering with two external dipoles (cf. Fig. 2.c) can measure both dipoles produced after a splitting, and thus probes the *dipole number fluctuations* in the target.

But the dipole picture breaks down at high density⁹, and thus cannot be used¹³ to also describe the $4 \rightarrow 2$ recombination process in Fig. 1.d. However, by viewing the diagram in Fig. 1.d upside down, it becomes clear that a merging process in the *target* can be alternatively interpreted as a splitting process in the *projectile*; since the latter is dilute, the dipole picture can then be used to describe its evolution. The two diagrams in Figs. 3.b and c represent the same processes as previously shown in Figs. 1.b and d, respectively, but now from the perspective of projectile evolution. In particular, the diagram in Fig. 3.c describes the simultaneous scattering of both child dipoles in the projectile, which is the unitarization mechanism at high energy. We see that the *unitarity corrections* correspond to either *gluon saturation*, cf. Fig. 1.d, or to *multiple scattering*, cf. Fig. 3.c, according to the frame in which we choose to view the evolution.

The previous considerations show that, in order to construct evolution equations which include both splitting and merging at large N_c , it is in fact sufficient to know the elementary *dipole splitting vertex*, that is, the probability density for an original color dipole to split into two new dipoles per unit rapidity. This has been first computed by Al Mueller⁹, and reads

$$\frac{dP}{d\tau} = \frac{\alpha_s N_c}{2\pi^2} \mathcal{M}(\mathbf{x}, \mathbf{y}, \mathbf{z}) d^2 \mathbf{z}, \quad \mathcal{M}(\mathbf{x}, \mathbf{y}, \mathbf{z}) \equiv \frac{(\mathbf{x} - \mathbf{y})^2}{(\mathbf{x} - \mathbf{z})^2 (\mathbf{y} - \mathbf{z})^2}, \quad (1)$$

where $d\tau$ is the rapidity increment ($\tau \sim \ln s$ is the rapidity gap between the projectile and the target), the pair $(\mathbf{x}, \mathbf{y})$ denotes the transverse coordinates of the quark and the antiquark in the original dipole, and $(\mathbf{x}, \mathbf{z})$ and $(\mathbf{z}, \mathbf{y})$ refer similarly to the two final dipoles. The latter have a common leg at $\mathbf{z}$, which physically represents the transverse coordinate of the emitted gluon.

By using Eq. (1) together with the picture of *projectile evolution*, cf. Fig. 3, one can immediately deduce the following evolution equation for the average scattering amplitude $\langle T(\mathbf{x}, \mathbf{y}) \rangle_\tau$




Figure 3: *Projectile evolution in the dipole picture.*

for a projectile made with a single dipole $(\mathbf{x}, \mathbf{y})$ (with $\bar{\alpha}_s = \alpha_s N_c / \pi$)¹⁴:

$$\frac{\partial}{\partial \tau} \langle T(\mathbf{x}, \mathbf{y}) \rangle_\tau = \frac{\bar{\alpha}_s}{2\pi} \int_{\mathbf{z}} \mathcal{M}(\mathbf{x}, \mathbf{y}, \mathbf{z}) \left\langle -T(\mathbf{x}, \mathbf{y}) + T(\mathbf{x}, \mathbf{z}) + T(\mathbf{z}, \mathbf{y}) - \langle T(\mathbf{x}, \mathbf{z}) T(\mathbf{z}, \mathbf{y}) \rangle_\tau \right\rangle. \quad (2)$$

Together, the first three terms in the r.h.s., which are linear in $\langle T \rangle$, represent the BFKL evolution, that is, the scattering of only one of the two child dipoles (cf. Fig. 3.b) and the reduction in the probability that the incoming dipole remains in its original state (the ‘virtual term’ $-\langle T(\mathbf{x}, \mathbf{y}) \rangle$). The last, non-linear, term $\langle T(\mathbf{x}, \mathbf{z}) T(\mathbf{z}, \mathbf{y}) \rangle_\tau$ describes the simultaneous scattering of both child dipoles, cf. Fig. 3.c. So long as the scattering is weak, $\langle T \rangle_\tau \ll 1$, this last term is negligible and the amplitude rises very fast (exponentially in τ), according to the BFKL equation. But when $\langle T \rangle_\tau \sim 1$, the non-linear term becomes important and tames the growth. Thus, as anticipated, the multiple scattering is the mechanism leading to unitarization.

But in order to study this mechanism in detail, one also needs the equation satisfied by the 2-dipole amplitude $\langle T(\mathbf{x}, \mathbf{z}) T(\mathbf{z}, \mathbf{y}) \rangle_\tau$, or, more generally, $\langle T(\mathbf{x}_1, \mathbf{y}_1) T(\mathbf{x}_2, \mathbf{y}_2) \rangle_\tau$. This involves, first, an evolution similar to Eq. (2) for any of the two incoming dipoles $(\mathbf{x}_1, \mathbf{y}_1)$ or $(\mathbf{x}_2, \mathbf{y}_2)$ (while the other one is simply a spectator). This will contribute terms linear in $\langle TT \rangle_\tau$ (the BFKL terms) together with unitarity corrections proportional to $\langle TTT \rangle_\tau$. Besides, the two external dipoles can ‘see’ a fluctuation in the gluon in the target, that is, they can directly probe the $2 \rightarrow 4$ splitting vertex in Fig. 1.c. To estimate this additional effect, it is preferable to adopt the point of view of *target evolution*, cf. Fig. 2.c, which implies³

$$\left. \frac{\partial \langle T(\mathbf{x}_1, \mathbf{y}_1) T(\mathbf{x}_2, \mathbf{y}_2) \rangle_\tau}{\partial \tau} \right|_{\text{fluct}} = \left(\frac{\alpha_s}{2\pi} \right)^2 \frac{\bar{\alpha}_s}{2\pi} \int_{\mathbf{u}, \mathbf{v}, \mathbf{z}} \mathcal{M}(\mathbf{u}, \mathbf{v}, \mathbf{z}) \mathcal{A}_0(\mathbf{x}_1, \mathbf{y}_1 | \mathbf{u}, \mathbf{z}) \mathcal{A}_0(\mathbf{x}_2, \mathbf{y}_2 | \mathbf{z}, \mathbf{v}) \times \nabla_{\mathbf{u}}^2 \nabla_{\mathbf{v}}^2 \langle T(\mathbf{u}, \mathbf{v}) \rangle_\tau. \quad (3)$$

The r.h.s. of this equation should be read as follows: A dipole $(\mathbf{u}, \mathbf{v})$ from the target splits into two new dipoles $(\mathbf{u}, \mathbf{z})$ and $(\mathbf{z}, \mathbf{v})$ with probability $(\bar{\alpha}_s/2\pi) \mathcal{M}(\mathbf{u}, \mathbf{v}, \mathbf{z})$ (cf. Eq. (1)), then the two child dipoles scatter off the external dipoles, with an amplitude^b $\alpha_s^2 \mathcal{A}_0$ for each scattering. Finally, the ‘amputated’ amplitude $(1/\alpha_s^2) \nabla_{\mathbf{u}}^2 \nabla_{\mathbf{v}}^2 \langle T(\mathbf{u}, \mathbf{v}) \rangle_\tau$ is, up to a normalization factor, the average dipole number density $\langle n(\mathbf{u}, \mathbf{v}) \rangle_\tau$ in the target.

Note that, although suppressed by an explicit factor α_s^2 , the contribution of the dipole number fluctuations to the evolution of $\langle TT \rangle_\tau$ is in fact a *dominant* effect in the *very dilute* regime where $\langle T \rangle_\tau \lesssim \alpha_s^2$: Indeed, in this regime $\langle T \rangle \langle T \rangle < \alpha_s^2 \langle T \rangle$, hence the dominant contribution to the rise of $\langle TT \rangle_\tau$ comes from fluctuations, via Eq. (3). This conclusion is in agreement with the qualitative analysis of the fluctuations in the companion article, Ref. ².

It is now clear that the evolution equations for dipole scattering amplitudes form an hierarchy, in which the general equation reads as follows (with $t = \bar{\alpha}_s \tau$ and $T^{(n)} \equiv \langle T(1) T(2) \dots T(n) \rangle_\tau$)

$$\frac{dT^{(n)}}{dt} = T^{(n)} - T^{(n+1)} + \alpha_s^2 T^{(n-1)}, \quad (4)$$

^b $\alpha_s^2 \mathcal{A}_0(\mathbf{x}, \mathbf{y} | \mathbf{u}, \mathbf{v})$ is the standard amplitude for dipole-dipole scattering via two-gluon exchange³.

in schematic notations which ignore combinatoric factors and the non-locality of the various vertices in the transverse space. The first two terms in the r.h.s. correspond to the BFKL evolution and the unitarity corrections, respectively, while the last term describes dipole number fluctuations in the target wavefunction. The iterative solution to these equations can be given a diagrammatic representation in terms of ‘BFKL Pomerons’ (the Green’s function of the BFKL equation) which split and merge with each other, and thus form *Pomeron loops*⁶.

An hierarchy like that in Eq. (4) can be exactly represented by a Langevin equation with a specific noise term. In statistical physics, this equation is known as the *stochastic FKPP equation* (sFKPP), and has the following generic structure (see Ref. ³ for details) :

$$\frac{dT}{dt} = T - T^2 + \sqrt{\alpha_s^2 T} \nu(t) \quad \text{with} \quad \langle \nu(t) \nu(t') \rangle = \delta(t - t'). \quad (5)$$

Such an equation represents a convenient starting point for numerical simulations, and particular limits of it have been already studied in this way¹⁵.

We shall conclude this brief presentation with a qualitative explanation of the concept of *self-duality* in the context of the high-energy QCD evolution^{5–7}. The seemingly trivial fact that a same $n \rightarrow 2$ vertex can be viewed as either merging in the target or splitting in the projectile has the physically important consequence that, in order to be *boost invariant*, the evolution equations for the scattering amplitudes must include the effects of *both* splitting and merging. Indeed, physical quantities must come out the same whatever is the system that we choose to evolve (the projectile or the target), since such a choice is tantamount to using a particular Lorentz frame. Let’s then assume that the *evolution Hamiltonian* H — the operator which governs the one-step evolution of the gluon fields in a hadronic wavefunction — contains a particular *merging* vertex $n \rightarrow 2$. When acting on the gluon fields in the *target*, this particular term in H generates an evolution for the scattering amplitudes which is equivalent to that that would be produced by a specific *splitting* vertex (the mirror, or ‘dual’, image of the original merging vertex) acting on the gluons in the *projectile*. Hence, in order to yield the same physical result when directly applied to the projectile, the Hamiltonian must also include the specific splitting vertex mentioned above, and therefore be *self-dual*. The precise duality transformation relating merging to splitting vertices depends upon the detailed structure of the scattering operator, and can be found in Refs.^{5–7}. The Hamiltonian underlying the evolution equations with Pomeron loops presented above is indeed self-dual⁶, and so is also the more general Hamiltonian in Refs.⁷.

1. For a review paper, see E. Iancu and R. Venugopalan, arXiv:hep-ph/0303204.
2. E. Iancu, these proceedings, arXiv:hep-ph/0510264.
3. E. Iancu, D. Triantafyllopoulos, *Nucl.Phys.* **A756**(2005)419; *Phys.Lett.* **B610**(2005)253.
4. For a recent review and more references, see C. Ewerz, arXiv:hep-ph/0306137.
5. A. Kovner, M. Lublinsky, *Phys.Rev.* **D71**(2005)085004; *Phys.Rev.Lett.* **94**(2005)181603.
6. J.-P. Blaizot, E. Iancu, K. Itakura, D.N. Triantafyllopoulos, *Phys. Lett.* **B615** (2005) 221.
7. Y. Hatta et al, arXiv:hep-ph/0504182; I. Balitsky, arXiv:hep-ph/0507237.
8. A.H. Mueller, A.I. Shoshi, S.M.H. Wong, *Nucl. Phys.* **B715** (2005) 440.
9. A.H. Mueller, *Nucl. Phys.* **B415** (1994) 373; *ibid.* **B437** (1995) 107.
10. E. Iancu and A.H. Mueller, *Nucl. Phys.* **A730** (2004) 460.
11. E. Levin, M. Lublinsky, arXiv:hep-ph/0501173; E. Levin, arXiv:hep-ph/0502243.
12. M. Braun, *Phys. Lett.* **B483** (2000) 115; arXiv:hep-ph/0504002.
13. E. Iancu, G. Soyez and D.N. Triantafyllopoulos, arXiv:hep-ph/0510094.
14. I. Balitsky, *Nucl.Phys.* **B463** (1996) 99; Yu.V. Kovchegov, *Phys. Rev.* **D60** (1999) 034008.
15. G. Soyez, *Phys. Rev.* **D72** (2005) 016007; R. Enberg et al, arXiv:hep-ph/0505101.